

# BOUNDARY CURRENTS AND HAMILTONIAN QUANTIZATION OF FERMIONS IN BACKGROUND FIELDS

JOUKO MICKELSSON

Theoretical Physics, Royal Institute of Technology, SE-10044,  
Stockholm, Sweden. e-mail jouko@theophys.kth.se

**ABSTRACT** The current algebra generated by fermions coupled to external gauge potentials and metrics on a manifold with boundary is discussed. It is shown that the previous methods, based on index theory arguments and used in the case without boundaries, carry over to the present problem. The resulting current algebra is the same as obtained from a quantization of bosonic Chern-Simons theories on space-times with nonempty boundaries. This means that also in the fermionic setting the construction is 'holographic', the Schwinger terms reside on the boundary.

The current algebra on manifolds with boundary has drawn recently great attention because of the correspondence between gravitation theories on anti-de Sitter type (odd dimensional) space-times and conformal field theory on the even dimensional boundary, [1,2]. In hamiltonian formulation one tries to construct the theory from the current algebra (for fixed time) on the odd dimensional boundary of the even dimensional space. In presence of gauge symmetries a prototype for the current algebra is an affine Lie algebra on the boundary in the case of a  $2+1$  dimensional interior space-time. This arises for example if the action functional (in the bulk) contains Yang-Mills and Chern-Simons terms, [4]. The boundary currents play also an important role in discussions of the quantum Hall effect, [3].

The purpose of the present letter is to show how the boundary current algebra arises from the canonical hamiltonian quantization of fermions in background gauge

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

fields. The method is closely related to previous studies of quantization of fermions in background gauge fields when the physical space-time is even dimensional and without boundary, [5,6], using a mixture of index theory and a variant of theory of gerbes.

Consider first an even dimensional compact Riemannian manifold  $M$  without boundary. We assume that  $M$  is oriented and with a fixed spin structure. Let  $V$  be a trivial vector bundle over  $M$  with an unitary action of a gauge group  $G$ . Let  $S$  be the tensor product of  $V$  and the Dirac spinor bundle over  $M$ . Given a gauge potential  $A$  in  $V$  and a metric  $g$  on  $M$  we have a Dirac operator  $D_{g,A}$  acting on the smooth sections of  $S$ . The Dirac operator on a compact manifold is Fredholm. We denote by  $H = H(g, A)$  the Hilbert space of square integrable sections of the bundle  $S$ . We denote by  $\mathcal{A}$  the space of all smooth vector potentials (with values in the Lie algebra  $\mathfrak{g}$  of the gauge group) and by  $\mathcal{M}$  the space of Riemann metrics on  $M$ .

We recall the construction of the Dirac determinant bundle (for a more detailed discussion and references to the extensive original literature, see [7, Chapter 9]). Let  $\lambda \in \mathbb{R}$  and  $U_\lambda$  the set of potentials and metrics such that  $\lambda$  is not in the spectrum of  $D_{g,A}$ . Then  $U_\lambda \subset \mathcal{M} \times \mathcal{A} = B$  is an open set and over  $U_\lambda$  we define a complex line bundle  $DET_\lambda$  such that the line at  $z \in U_\lambda$  is

$$(1) \quad DET_\lambda(z) = \wedge^{top}(S_\lambda^+) \otimes \wedge^{top}(S_\lambda^-)^*,$$

where  $S_\lambda^\pm$  is the space of positive (negative) chirality modes in the spectral subspace  $(D_z)^2 < \lambda$ .

Let  $0 < \lambda < \lambda'$ . We define  $DET_{\lambda\lambda'}$  to be a trivial complex line bundle over the intersection  $U_{\lambda\lambda'} = U_\lambda \cap U_{\lambda'}$ . The line at  $z$  is the tensor product

$$(2) \quad DET_{\lambda\lambda'}(z) = \wedge^{top}(S_{\lambda\lambda'}^+) \otimes \wedge^{top}(S_{\lambda\lambda'}^-)^*,$$

where  $S_{\lambda\lambda'}^\pm = (S_\lambda^\pm)^\perp \cap S_{\lambda'}^\pm$ . Clearly  $DET_{\lambda'} = DET_{\lambda\lambda'} \otimes DET_\lambda$  over  $U_{\lambda\lambda'}$ . Furthermore,  $DET_{\lambda\lambda'}$  has a canonical trivialization defined as follows. Let  $v_1, v_2, \dots, v_n$  be any orthonormal basis of  $S_{\lambda\lambda'}^+$ . Then the vectors  $Jev_i$  form a basis of  $(S_{\lambda\lambda'}^-)^*$ , where  $\epsilon = D_z/|D_z|$  and  $J$  is the canonical antilinear isomorphism from a Hilbert space to its dual. Note here that  $D_z$  anticommutes with the chirality operator whereas  $|D_z|$  commutes with chirality and thus  $\epsilon$  maps  $S^+$  to  $S^-$ . As a consequence of eq.

2 and the triviality of the bundles  $DET_{\lambda\lambda'}$ , the local line bundles  $DET_\lambda$  patch up to a globally defined line bundle  $DET$  over  $B$ . The construction is gauge equivariant, leading to a line bundle  $DET$  over  $B/(Diff_0(M) \times \mathcal{G}_0)$ . Here  $\mathcal{G}_0$  is the group of based gauge transformations, i.e., those gauge transformations which at a fixed base point  $p \in M$  take the value 1.  $Diff_0(M)$  is the group of diffeomorphisms  $h$  such that  $h(p) = p$  and the derivative of  $h$  at  $p$  is the identity.

If the Dirac operator  $D_z$  does not have zero modes then the construction of the fermionic Fock space and the vacuum is canonical. The representation of the CAR algebra is fixed by the polarization  $H(z) = H_+(z) \oplus H_-(z)$  of the one-particle Hilbert space to positive and negative spectral subspaces. The vacuum is characterized by the property

$$a^*(u)|z\rangle = 0 = a(v)|z\rangle$$

for all  $u \in H_-$  and  $v \in H_+$ . The only nonzero anticommutators are  $[a^*(u), a(v)]_+ = \langle v, u \rangle$ , the inner product in  $H$  being antilinear in the first argument.

In the case there are zero modes there is an ambiguity. One has to decide which of the zero modes should belong to  $H_+$  and which to  $H_-$ . The choice cannot be made in an arbitrary manner if we want to have a smooth bundle of Fock spaces. We resolve this ambiguity by requiring that the vacuum is annihilated by all  $a^*(v)$ 's for positive chirality zero modes and all  $a(u)$ 's corresponding to negative chirality zero modes. That is, we may view the vacuum as an element in the line

$$DET = \wedge^{top}(S_{D_z=0}^+) \times \wedge^{top}(S_{D_z=0}^-)^*$$

belonging to the space  $\wedge(S_{D_z=0}^+) \otimes \wedge(S_{D_z=0}^-)^*$ .

To be more precise and in order to show the smoothness of this construction one can proceed as before in the case of the bundle  $DET$ . First define local Fock bundles  $\mathcal{F}_\lambda$  over  $U_\lambda$ , with  $\lambda > 0$ , as follows. Let  $H_{+,\lambda}$  (resp.  $H_{-,\lambda}$ ) be the spectral subspace  $D_z < \lambda$  (resp.  $D_z < -\lambda$ ). Set

$$(3) \quad \mathcal{F}_\lambda(z) = \wedge(H_{+,\lambda} \oplus S_\lambda^+) \otimes \wedge(H_{-,\lambda} \oplus S_\lambda^-)^*.$$

The vacuum  $|z, \lambda\rangle$  in  $\mathcal{F}_\lambda(z)$  is defined as a normalized element in  $DET_\lambda(z) \subset \mathcal{F}_\lambda(z)$ . Wedging this with the canonical element in  $DET_{\lambda\lambda'}$  we obtain an element in  $DET_{\lambda\lambda'}(z) \subset \mathcal{F}_{\lambda'}(z)$ . This provides us an identification of  $\mathcal{F}_\lambda$  with  $\mathcal{F}_{\lambda'}$ . Thus we obtain a globally defined bundle of Fock spaces  $\mathcal{F}$  over  $B$ .

The creation and annihilation operators are defined in a Fock space  $\mathcal{F} = \wedge(W) \otimes \wedge(W^\perp)^*$  in the standard way. The creation operator  $a^*(u)$  for  $u \in W$  is represented by the exterior product  $u \wedge$  and the annihilation by the contraction  $i_u$  (which is antilinear in  $u$ ). The creation and annihilation operators for  $v \in W^\perp$  are  $a^*(v) = i_{Jv}$  and the annihilation operator  $a(v) = Jv \wedge$ .

In the case of a manifold with boundary there are important modifications to the construction above. We have to specify elliptic boundary conditions for the Dirac operators  $D_z$ .

We first consider the APS boundary conditions, [8], defined by  $\psi|_{\partial M} \in P_\lambda(H^b)$  over those parameters  $z = (g, A)$  such that  $\lambda \notin \text{Spec}(h_z)$ . Here  $P_\lambda$  is the spectral projection to the part of the boundary Hilbert space  $H^b$  where  $h_z < \lambda$ ;  $h_z$  is the boundary Dirac operator to be defined below. In order to apply the APS theory we have to assume that the geometry is tubular near the boundary. This means that the metric is a product metric near  $\partial M$  and the gauge potentials have vanishing normal derivatives at the boundary. The Dirac equation near  $\partial M$  can be written as

$$\frac{\partial}{\partial t} \psi = h_z \psi$$

where  $t$  is a coordinate in the normal direction near the boundary,  $h_z$  is the (hermitean) boundary operator  $\gamma_t \gamma^i \nabla_i$ ,  $i = 1, 2, \dots, 2n-1$  and  $\nabla_i$  denotes the combined gauge and Levi-Civita covariant derivative in the  $i$ :th local basis vector direction.

The operator  $h_z$  commutes with the chirality operator  $\gamma_{2n+1}$  and therefore the positive and negative eigenvalues for the Dirac operator in the bulk come in pairs like in the case without boundary and the determinant line bundle and the Fock spaces are defined as before. The crucial difference is in the fact that the determinant line bundle (and therefore also the Fock spaces) are only defined locally, over the sets  $V_\lambda = \{z \in B | \lambda \notin \text{Spec}(h_z)\}$ , in the parameter space.

In order to define the Fock spaces globally one can choose a spectral section on the boundary (for an alternative construction see [6]). We denote by  $DET(D, P)$  the determinant bundle for the family  $D_z$  of Dirac operators with the boundary conditions  $\psi|_{\partial M} \in P(H^b(z))$ . Here  $P = P^{(z)}$  is a projection operator acting in the boundary Hilbert space  $H^b(z)$ . It should depend smoothly on the parameter  $z$  and the difference  $P - P_\lambda$  must be Hilbert-Schmidt. The determinant bundle in the

bulk becomes then

$$(4) \quad DET(D, P) = DET(D, P_\lambda) \otimes DET_F(P_\lambda^+, P^+)$$

where  $P$  is the spectral section, [10]. We have set  $P^+ = \pi_+ P$  where  $\pi_\pm$  are the chiral projectors on the boundary. Since  $P^{(z)}$  and  $P_\lambda(z)$  are Hilbert-Schmidt related, the latter line bundle  $DET(P_\lambda^+, P^+)$  can be taken as the (relative) Fredholm determinant bundle over a restricted Grassmannian manifold as discussed in [11]. We have also performed the standard identification of  $DET(D)$  as the tensor product  $\wedge^{top}(ker D^+) \otimes \wedge^{top}(coker D^+)^*$  which leads to the localization of the domain to the positive chirality sector in (4).

The gauge and diffeomorphism group action on the first factor in the above formula (4) is equivariant but not on the second factor because in general there is no smooth equivariant choice for the spectral section  $P$ . It follows that there is a potential obstruction to lifting the group actions to the total space of the Fock bundle due to a nontrivial extension coming from the lifting of the group actions on potentials and metrics to the complex line bundle  $DET_F(P_\lambda^+, P^+)$ . On the level of Lie algebras this obstruction is seen as nontrivial Schwinger terms in current commutation relations. As explained in [6] the Schwinger terms can be computed from the curvature of the determinant bundles. APS index theory determines the cohomology class of the curvature form.

Any parameter  $z \in B$  determines a boundary geometry  $z_b$ . On the other hand, for any  $z_b$  we can choose a continuation to a metric and gauge connection  $\hat{z}$  in the bulk. The map  $z \mapsto \hat{z}$  clearly defines a map from  $B/(Diff_0 \times \mathcal{G}_0)$  to a contractible set of parameters  $\hat{z}$  (since the space of the parameters  $z_b$  is contractible). Let  $1 - P^{(z)}$  be the *Calderon projection* associated to the Dirac operator  $D_{\hat{z}}$ . By Calderon projection we mean the projection to the space of boundary values of solutions of the Dirac equation  $D_{\hat{z}}\psi = 0$ . It is known that  $P^{(z)}$  differs by a Hilbert-Schmidt operator (in fact, by a smoothing operator) from the spectral projection  $P_\lambda$  for the boundary hamiltonian; if we drop the requirement concerning the vanishing of the normal derivatives of metrics and potentials at the boundary, one has more general  $L_p$  type estimates replacing the Hilbert-Schmidt condition, [12]. It follows that we may choose  $P_z$  as the spectral section above. Note that now everything in the parametrization of the determinant lines in  $DET(P_\lambda^+, P^+)$  depends on the

*boundary data only.*

The curvature of  $DET(P_\lambda^+, P^+)$  can be computed from the APS theory. For determining the Schwinger terms one needs to know the curvature to the gauge directions only, [6]. The result is equivalent to a standard local quantity; when the boundary is 1-dimensional one gets the central term in an affine Lie algebra and in the 3-dimensional case it is extension part in the Mickelsson-Faddeev algebra.

Following [9] we can construct explicitly the vacuum bundle, that is, the chiral Dirac determinant bundle from the chiral anomaly functional. Restricting to the case of a fixed metric and  $\partial M = \emptyset$  the sections of the bundle  $DET$  are complex functions  $\psi : \mathcal{A} \rightarrow \mathbb{C}$  such that

$$(5) \quad \psi(A^g) = \psi(A) \cdot e^{i\omega(g;A)}$$

where  $A^g = g^{-1}Ag + g^{-1}dg$  and the anomaly functional  $\omega$  is an integral of a local differential polynomial in  $A$  and  $dg g^{-1}$ . It satisfies the cocycle property

$$(6) \quad \omega(g_1 g_2; A) = \omega(g_1; A) + \omega(g_2; A^{g_1}).$$

The important property of this construction is that it extends to the case with boundary. The condition (5) should now hold for all potentials  $A$  which obey the tubularity condition at the boundary and for those gauge transformations  $g$  which smoothly approach the value 1 at the boundary. Let  $\mathcal{G}_{00}$  be the group of gauge transformations with vanishing normal derivatives at the boundary and which are equal to 1 on the boundary. We can mod out by  $\mathcal{G}_{00}$  and we obtain a determinant bundle over  $\mathcal{A}/\mathcal{G}_{00}$ . We have not specified explicitly the boundary conditions  $P$  as a function of the boundary data. The reason is that *whatever choice we make for the boundary projections  $P$  (satisfying the Hilbert-Schmidt relations and depending smoothly on boundary data) the resulting bundles will be equivalent.* This is seen from (4). If we change the boundary conditions then the corresponding determinant bundles are related by the factor  $DET(P^+, P'^+)$ . Modding out by the gauge and diffeomorphism symmetries in the bulk does not affect the boundary data. So even after pushing things to  $\mathcal{A}/\mathcal{G}_{00}$  the bundle  $DET(P^+, P'^+)$  will be a bundle over the affine space of all boundary potentials and therefore automatically trivial. Clearly, the same comments apply to metrics and diffeomorphisms.

The bigger group  $Map(M, G)$  of all gauge transformations (including also those which are not equal to 1 at the boundary but still have vanishing normal derivatives) acts projectively in the space of sections  $\Gamma(DET)$  through the formula

$$(T(g)\psi)(A) = e^{i\gamma(g;A)}\psi(A^g),$$

where  $\gamma$  is an integral of another local differential polynomial, [9]. This functional does not satisfy the cocycle condition. This means that  $T(g_1)T(g_2) = T(g_1g_2)e^{i\omega_2(g_1, g_2; A)}$  for some 2-cocycle  $\omega_2$ . Infinitesimally this means that the Lie algebra of  $Map(M, G)$  is extended by an abelian ideal consisting of functions  $f : \mathcal{A} \rightarrow \mathbb{C}$ ,

$$(7) \quad [(X, f), (Y, g)] = ([X, Y], X \cdot g - Y \cdot f + c(X, Y; A)),$$

where  $c$  is a Lie algebra cocycle and the action  $X \cdot g$  is the Lie derivative of a function determined by the infinitesimal gauge transformation  $\delta_X A = [A, X] + dX$ .

As promised in the abstract, a basic property of the functional  $c(X, Y; A)$  is that it depends only on the boundary values of the fields, [9]. In  $2 + 1$  dimensions we have

$$(8) \quad c(X, Y; A) = \frac{i}{2\pi} \int_{M_2} \text{tr} dX \wedge dY = \frac{i}{2\pi} \int_{\partial M_2} \text{tr} X dY,$$

where  $M_2$  is the 2-dimensional physical space. In  $4 + 1$  dimensions we have

$$(9) \quad \begin{aligned} c(X, Y; A) &= \frac{i}{24\pi^2} \int_{M_4} \text{tr} dA \wedge (dX \wedge dY - dY \wedge dX) \\ &= \frac{i}{24\pi^2} \int_{\partial M_4} \text{tr} A \wedge (dX \wedge dY - dY \wedge dX). \end{aligned}$$

## References

- [1] J. Maldacena: The large  $N$  limit of superconformal field theories and supergravities. *Adv. Theor. Math. Phys.* **2**, 231 (1998). hep-th/9711200
- [2] E. Witten: Anti-de Sitter space and holography. *Adv. Theor. Math. Phys.* **2**, 253 (1998). hep-th/9802150
- [3] J. Fröhlich and U.M. Studer: Gauge invariance and current algebra in non-relativistic many-body theory. *Rev. Modern Phys.* **65**, no. 3, part 1, 753 (1993)
- [4] J. Mickelsson: On a relation between massive Yang-Mills theories and dual string models. *Lett. Math. Phys.* **7**, 45 (1983)
- [5] J. Mickelsson: On the hamiltonian approach to commutator anomalies in  $3 + 1$  dimensions. *Phys. Lett.* **B241**, 70 (1990)
- [6] A. Carey, J. Mickelsson, and M. Murray: Index theory, gerbes, and hamiltonian quantization. *Commun. Math. Phys.* **183**, 707 (1997); Bundle gerbes applied to quantum field theory. hep-th/9711133, to be publ. in *Rev. Math. Phys.*
- [7] N. Berline, E. Getzler, and M. Vergne: *Heat Kernels and Dirac Operators*. Grundlehren der mathematischen Wissenschaften 298. Springer-Verlag (1992)
- [8] M.F. Atiyah, V.K. Patodi, and I.M. Singer: Spectral asymmetry and Riemannian geometry I-III. *Math. Proc. Camb. Phil. Soc.* **77**, 43 (1975); **78**, 405 (1975); **79**, 71 (1976).
- [9] J. Mickelsson: Kac-Moody groups, topology of the Dirac determinant bundle and fermionization. *Commun. Math. Phys.* **110**, 173 (1987). *Current Algebras and Groups*. Plenum Press, London and New York (1989)
- [10] S. Scott: Splitting the curvature of the determinant line bundle. math.AP/9812124. *Proc. Amer. Math. Soc.* (to appear)
- [11] A. Pressley and G. Segal: *Loop Groups*. Clarendon Press, Oxford (1986)
- [12] L. Fiedlander and A. Schwarz: Grassmannian and elliptic operators. *funct-anal/9704003*